Mon. Not. R. Astron. Soc. **000**, 1–10 (2007) Printed 12 August 2008 (MN LATEX style file v2.2)

Proper motions of field L and T dwarfs -II

S. L. Casewell[1⋆], R. F. Jameson[1] and M. R. Burleigh[1]
[1]*Department of Physics and Astronomy, Univeristy of Leicester, Univeristy Road, Leicester, LE1 7RH, UK*

arXiv:0808.1633v1 [astro-ph] 12 Aug 2008

ABSTRACT
By using images taken with WFCAM on UKIRT and SofI on the NTT and combining them with 2MASS we have measured proper motions for 126 L and T dwarfs in the dwarf archive. Two of these L dwarfs appear to have M dwarf common proper motion companions, and 2 also appear to be high velocity dwarfs, indicating possible membership of the thick disc. We have also compared the motion of these 126 objects to that of numerous moving groups, and have identified new members of the Hyades, Ursa Major and Pleiades moving groups. These new objects, as well as those identified in Jameson et al. (2008a) have allowed us to refine the L dwarf sequence for Ursa Major that was defined by Jameson et al. (2008b).

Key words: stars:kinematics-stars:low-mass,brown dwarfs - open clusters and associations:individual:Ursa Major:Hyades:Pleiades

1 INTRODUCTION

Brown dwarfs may be thought of as failed stars. These low mass ($\leqslant$70 M_{Jup} Burrows et al. 2001), cool objects are the lowest mass objects that the star formation process can produce. The majority of the brown dwarfs that have been discovered to date are field objects found using surveys such as the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006, see Leggett et al. 2002 for examples), the DEep Near-Infrared Sky survey (DENIS; DENIS Consortium 2005, see Delfosse et al., 1999 for examples), the Sloan Digital Sky Survey (SDSS;York et al. 2000 see Hawley et al. 2002 for examples) and the UKIRT Deep Infrared Sky Survey (UKIDSS; Lawrence et al. 2007, see Kendall et al. 2007; Lodieu et al. 2007a for examples). However, to study brown dwarfs in depth, a knowledge of their age is essential, which means we must study brown dwarfs in open star clusters or moving groups.

Once a brown dwarf has been proved to belong to an open star cluster, or a moving group, then its age is known, allowing meaningful comparisons to evolutionary models to be made. The most recent example of this is the study done by Bannister & Jameson (2007) who used existing proper motions and parallax measurements to show that a selection of field dwarfs in fact belong to the Ursa Major and Hyades moving groups. The importance of this study, is that these are the first brown dwarfs to be associated with an older cluster or group (age $>$200 Myr). Older clusters such as the Hyades are expected to contain very few or no brown dwarfs or low mass members, due to the dynamical evolution of the cluster over time (Adams et al. 2002). However, these escaped low mass objects may remain members of the much larger moving group that surrounds the cluster.

Jameson et al. (2008a) followed this work by using the wide field camera (WFCAM, Casali et al. 2007) on the United Kingdom Infrared Telescope (UKIRT) to image 143 known field L dwarfs. These images provided a second epoch for proper motion measurements, when combined with existing 2MASS images, typically taken 7 years previously. Using the proper motions and a distance calculated using the spectral type of the L dwarf given by Cruz et al. (2003), the moving group method was applied, and all 143 objects were scrutinised to check if their direction and magnitude of motion made them candidates of the many moving groups known. Members of the Hyades, Ursa Major and Pleiades moving groups were found. Radial velocity measurements such as those of Zapatero Osorio et al. (2007) are required however, before it can be determined if these moving group members are cluster members that have "escaped" as the cluster has dynamically evolved (Adams et al. 2002). Is should also be noted that galactic resonances can produce effects similar to moving groups, and so all members may not be coeval (Dehnen 1998).

To continue the study started by Bannister & Jameson (2007) and Jameson et al. (2008a), we have measured proper motions for the majority of the remaining known field L dwarfs listed in the online L and T dwarf archive (http://spider.ipac.caltech.edu/staff/davy/ARCHIVE/). This has again been accomplished using the WFCAM on UKIRT and for the more southern objects, Son of ISAAC (SofI) on the 3.58m ESO New Technology Telescope (NTT). Using these wide field images and existing catalogue data, we have measured proper motions for an additional 126 L and T dwarfs listed in the dwarf archive.

These proper motion data may be put to a number of uses. Using reduced proper motion diagrams they can be used as an approximate measure of distance. The proper motion measurements can also be used to help identify objects as members of a star cluster or members of a moving group via the moving cluster method.

⋆ E-mail: slc25@star.le.ac.uk

Taken with measured radial velocities and distances, it can yield all three components of velocity (U,V,W). As brown dwarfs tend to be faint, measuring their radial velocities is very difficult. As a result, very few L or T dwarfs have measurements, none of which are in this sample. This means we cannot determine whether these moving group members are escaped cluster members or otherwise.

Our proper motion data are discussed and listed in section 2 of this paper.

2 PROPER MOTION MEASUREMENTS

2.1 Data acquisition and reduction

In order to measure proper motions for known L and T dwarfs we observed a sample of the known L and T dwarfs from the online dwarf archive (detailed in Tables 1 and 2, see http://spider.ipac.caltech.edu/staff/davy/ARCHIVE/ for discovery references) with *J* band magnitudes of less than 16.5. 143 of these objects were presented in Jameson et al. (2008a). To complete the sample, 126 additional objects have been observed, 88 with declinations between -30° and +60°, and 38 with declinations of less than -30°. This first group were imaged using WFCAM on UKIRT over the period of June 2006 to March 2008. WFCAM is a near infrared imager consisting of 4 Rockwell Hawaii-II (HgCdTe 2048x2048) arrays arranged such that 4 separately pointed observations can be tiled together to cover a filled square of sky covering 0.75 square degrees with 0.4 arcsecond pixels (Casali et al. 2007). However, as we only required the image of the brown dwarf in question, we only used array 3 which is regarded as the least noisy array. WFCAM is ideal for this work, as the large field of view per chip means there are many other stars in the image, which can be used as astrometric reference stars. The images were taken in the *J* band in non-photometric conditions using exposure times of ≈5-10 minutes and a nine point dither pattern. These exposure times gave S/N≈100 even in the poor conditions.

The images were reduced at the Cambridge Astronomical Survey Unit (CASU) using procedures which have been custom written for the treatment of WFCAM data (Irwin et al., in preparation, Dye et al. 2006.). In brief, each frame was debiased, dark corrected and then flat fielded. The individual dithered images were stacked before having an object detection routine run on them. The frames were astrometrically calibrated using point sources in the 2MASS catalogue. The accuracy is typically ≈0.1" (Dye et al. 2006) The photometric calibration employed by the CASU pipeline also relies on 2MASS data (there are typically hundreds of 2MASS calibrators per detector) and is found to be accurate to ≈2% in good conditions (Warren et al. 2007). However as we wished to measure proper motions, the astrometric calibration was more important than the photometric calibration for these data.

The objects with declinations less than -30° were observed using SofI on the NTT. SofI is a wide field infrared imager consisting of an Hawaii HgCdTe 1024x1024 array. This has a plate scale of 0.292 "/pixel and a field of view of ∼ 5 arcminutes (Moorwood, Cuby & Lidman 1998). The images were taken between 02/04/2007 and 31/08/2007 in the *J* band. The conditions were non-photometric, exposure times were between 5 and 10 minutes, and a nine point dither pattern was used. As for the WFCAM images, S/N≈100 was achieved in the conditions.

These data were reduced using the ORAC-DR pipeline (Allan et al. 2002). This pipeline is maintained and developed by the JAC in Hawaii, and has been adapted to reduce SofI data (Currie 2003). The pipeline bias corrected, flat fielded, sky subtracted and created mosaics out of the nine images per object. These mosaics were then astrometrically calibrated using point sources in 2MASS, and the STARLINK Autoastrom package. The accuracy of this calibration is typically greater than 0.1", as measured by the rms of the fit to the 2MASS point sources. This is worse than the WFCAM astrometry as one might expect, as there are much fewer reference stars in the smaller SofI images. Object detection was completed using SExtractor using the image parameters (gain etc) and a suitably sized background mesh, tailored to the seeing and pixel scale of the image.

2.2 Calculating proper motions

The astrometry for 2MASS is good to 80 mas over the whole survey, and to 50 mas over a small area (Skrutskie et al. 2006). Because the WFCAM and SofI astrometry is also calibrated to the 2MASS catalogue, accurate relative proper motion measurements could be calculated simply by taking the difference in 2MASS and WFCAM positions and dividing by the epoch difference. We calculated the epoch difference by taking the difference in the Julian date as given in the FITS header for each image, which is between 6.5 and 8.7 years with the average epoch difference being 7.8 years for SofI and between 5.8 and 9.5 with an average of 8.5 for WFCAM. The proper motion measurements for each object in every WFCAM array 3 and SofI image were calculated by this method. This same method was used for our previous 143 objects (Jameson et al. 2008a) and produced results consistent with the existing data.

This motion was then converted into mas yr^{-1}. The proper motion has been calculated directly from the RA and dec of the object in question, not from pixel motion on images, hence μ_α=($\Delta\alpha$/ΔT)cos δ " yr^{-1} if $\Delta\alpha$ is converted to arcseconds. These proper motions are relative proper motions, in the sense that they are relative to the bulk of the background stars in the field, which are generally moving slowly enough to be assumed to have zero motion.

However, for the WFCAM fields we checked the reference star motion so that the proper motion of the brown dwarf could be altered if there was a standard offset in the field.

The proper motions were separated in $\mu_\alpha cos\delta$ and μ_δ from -500 to 500 mas yr^{-1} in each direction, in bins of size 20 mas yr^{-1}, and the number of objects falling into each bin were totalled. We then fitted a two dimensional Gaussian to the data for each field to determine the spread of the reference stars, as well as the true centre of the motion. The process was then repeated after the initial fit, rejecting any objects that lay outside 3σ of the fitted Gaussian, before fitting another Gaussian to this data. This fitting was important in some cases as the reference stars had quite a large spread, and in other cases the proper motion of our brown dwarf was of the same order of magnitude as the references. These centroiding changes, or the centres of the fit, were then subtracted from the calculated proper motion measurements.

We used the σ value of the Gaussian to determine the error on our measurements. In general the errors were of the order of ≈ 15 mas yr^{-1}. The quoted errors are the σ value of the Gaussian fitted to the proper motion points for each image. Strictly this should be the σ plus the position error of the object added in quadrature. However, the centroiding errors for WFCAM are less than 2 mas yr^{-1}, and so are small compared to the σ value.

This method could not be used for some of the SofI fields, as there were less than 30 reference stars present, and in some cases, as few as 6. For these fields, the Gaussian fitting did not converge.

As an alternative method of gauging the offset in motion and the proper motion errors, the mean of both $\mu_\alpha \cos\delta$ and μ_δ was calculated for all of the objects in the field, once the brown dwarf had been removed. The standard deviation was also calculated, and again an iterative process used to remove any objects that lay further from the mean than 3 standard deviations. The process was then repeated and the means were then used as the offset in motion and the standard error of the mean values was then used as the error of the proper motion. The measured proper motions are given in Table 1 for the brown dwarfs imaged by SofI and in Table 2 for the brown dwarfs imaged by WFACM.

Some of the L dwarfs have had their proper motions measured by other people namely Caballero (2007) and Schmidt et al. (2007). The proper motions given by Caballero (2007) are taken from the SuperCOSMOS Sky Survey. The errors on the proper motions are typically 10 mas yr^{-1}, although in some cases this is larger, such as in the case of J0847-15 and J0909-07. This is because the SuperCOSMOS sky survey which consists of scanned sky atlas photographic plates, imaged in at least two epochs (Hambly et al. 2001a,b), has necessarily measured the proper motions over an unfavourable (i.e. short) epoch difference (using the minimum of 2 plates required to calculate the motions). Both of these objects have Schmidt et al. (2007) proper motions that do agree with our values, and so the Caballero (2007) measurements are taken to be inaccurate. Tables 3 and 4 show the proper motion comparisons as given by Caballero (2007) and Schmidt et al. (2007).

It should be pointed out that 0915+04 is described as a binary by Reid et al. (2006) with a separation of 0.73" at -155.0°. Our image also shows two resolved objects which may be why there is such a difference between the Schmidt et al. (2007) measurements and these data.

Two further two objects have additional proper motion measurements. J0255-47 has a proper motion measurement of 1.14±0.0022" yr^{-1} and a θ measurement of 119.5±0.21° (Costa et al. 2006). Our measurements are μ=1.18±0.01 " yr^{-1} and θ=117.4±1 °. This object also has an additional parallax measurement of $\sim$ 5 pc (Costa et al. 2006), placing it well within the local neighbourhood.

J0320-04 is suspected to be an unresolved M8.5+T5 binary (Burgasser et al. 2008) and has a proper motion of 0.562±0.005, and a θ of 205.9, which compares favourably with our measurements of 0.65±0.01" and 206.5±2 degrees.

POSSIBLE BINARIES

This study also provides an opportunity to search for wide common proper motion companions to the known brown dwarfs. Any objects within the field that have a proper motion within 15 mas yr^{-1} of the dwarf's motion were considered as possible companions. Obviously any brown dwarf with a proper motion close to zero, or the majority of background sources, will have a large number of "companions" within our selection. These objects have been excluded.

Two brown dwarfs J100-33 and J0147+34 appear to have common proper motion companions.

J1004-33 has a possible M dwarf companion which SIMBAD names as LHS5166, a high proper motion star with proper motion of $\mu_\alpha \cos\delta$=400 mas yr^{-1} and μ_δ=-420 mas yr^{-1}. Our measurement of its proper motion is 370.66±25.14, -342.62±14.46 mas yr^{-1}.

J0147+34 has a companion which is an M dwarf as identified by Wei et al. (1999). This M dwarf is an X-ray source (2E 0144.7+3438 as given by SIMBAD), suggesting that it may be young. It is 43" away from the known brown dwarf and has a proper motion of 46.2±18.3, -46.4±14.8 mas yr^{-1}.

Further proper motion and radial velocity measurements are required before these can be confirmed as true companions however.

FAST MOVING OBJECTS

In paper I (Jameson et al. 2008a), we discovered 8 high velocity L dwarfs which we suggested may belong tho the thick disk or halo population. Applying the same criteria to these data (that the proper motion must be greater than 0.85" yr^{-1}), we found 3 candidates from the SofI images. The distance estimate derived from their spectral type as defined in Cruz et al. (2003) is 22.35 pc for J0014-48 (L2), 5.0 pc for J0255-47 (L8) and 25.8 pc for J2150-75 (L1). J0255-47, also has a parallax measurement from Costa et al. (2006), which is 4.96 pc, in agreement with the spectral type distance. This object is relatively nearby, which gives it a true velocity of only 28 km s^{-1}. The other two objects have velocities of 95 km s^{-1} and 112 km s^{-1} respectively. Both J0014-48 and J2150-75 have relatively blue $J - K_S$ and $H - K_S$ colours for their spectral types, as with other high velocity dwarfs (Jameson et al. 2008a; Schmidt et al. 2007), indicating that they too may be members of an older ($\sim$ 10 Gyr), thick disk population.

From the WFCAM data, just one object met the required selection criteria, J0921-21, an L2 dwarf, with a spectral type distance of 12.4 pc, and a velocity of 55 km s^{-1}. This velocity is not high enough for this object to be considered a true high velocity dwarf.

MOVING GROUPS

We have attempted to determine if any of these brown dwarfs are members of known moving groups. Being a moving group member means that an age can be estimated for the dwarf. This does not necessarily mean that the moving group member is an "escaped" cluster member however, as galactic resonances can produce a similar effect (Dehnen 1998). To determine if any of these field dwarf are moving group members, we used the moving cluster method as described in Bannister & Jameson (2007) and Jameson et al. (2008a). Again, we used the proper motion measurements and direction of that motion as well as a distance derived from the spectral type as described in Cruz et al. (2003). If the difference in angles between the measured and predicted motion towards the convergent point is less than 14° plus the error on the measurement, and the ratio between the moving cluster distance and the Cruz spectral type distance was greater than 0.72 and less than 1.28 as in Bannister & Jameson (2007) and Jameson et al. (2008a), then the selected objects were plotted on a colour−magnitude diagram, and compared to the DUSTY and NextGen models (Chabrier et al. 2000; Baraffe et al. 1998) and to empirical isochrones defined by known cluster members (Casewell et al. 2007 for the Pleiades and Hogan et al. 2008 for the Hyades.)

Hyades

The Hyades cluster has a distance of 46 pc and covers $\approx 20°$ of the sky. The Hyades has, until recently, been thought to contain almost no low mass members. Extensive searches such as those

Table 1. 2MASS name, RA, Dec, $\mu_\alpha \cos\delta$, μ_δ and 2MASS magnitudes for all the L and T dwarfs for which we measured proper motions using SofI.

Name 2MASS	RA J2000	Dec J2000	$\mu_\alpha \cos\delta$ mas yr^{-1}	μ_δ mas yr^{-1}	J	H	K_S
J00145575-4844171	00 14 55.76	-48 44 17.15	851.29 ± 12.38	279.42±8.02	14.050± 0.033	13.107± 0.035	12.723± 0.028
J00165953-4056541	00 16 59.53	-40 56 54.15	162.51 ± 14.44	16.06±5.38	15.316± 0.060	14.206± 0.047	13.432± 0.037
J00325584-4405058	00 32 55.84	-44 05 05.87	116.76 ± 7.85	-88.96±5.45	14.776± 0.032	13.857± 0.032	13.269± 0.035
J00531899-3631102	00 53 18.99	-36 31 10.29	36.82 ± 22.71	-72.41±8.57	14.445± 0.023	13.480± 0.030	12.937± 0.027
J01174748-3403258	01 17 47.48	-34 03 25.82	103.14 ± 13.98	-39.70±7.00	15.178± 0.034	14.209± 0.038	13.489± 0.036
J01253689-3435049	01 25 36.90	-34 35 04.91	120.42 ± 42.95	-12.97±21.05	15.522± 0.054	14.474± 0.051	13.898± 0.054
J01415823-4633574	01 41 58.23	-46 33 57.43	92.90 ± 10.43	-5.54±8.27	14.832± 0.041	13.875± 0.024	13.097± 0.030
J02182913-3133230	02 18 29.13	-31 33 23.08	-131.09 ± 9.87	-97.18±16.69	14.728± 0.038	13.808± 0.036	13.154± 0.033
J02550357-4700509	02 55 03.58	-47 00 50.99	1052.88 ± 11.17	-546.54±6.17	13.246± 0.024	12.204± 0.022	11.558± 0.023
J03185403-3421292	03 18 54.04	-34 21 29.22	401.63 ± 9.93	43.28±4.10	15.569± 0.053	14.346± 0.043	13.507± 0.038
J03572695-4417305	03 57 26.96	-44 17 30.55	63.15 ± 12.71	2.35±7.54	14.367± 0.029	13.531± 0.025	12.907± 0.026
J04430581-3202090	04 43 05.81	-32 02 09.01	-0.46 ± 13.70	205.19±7.54	15.273± 0.050	14.350± 0.055	13.877± 0.062
J04455387-3048204	04 45 53.88	-30 48 20.46	158.07 ± 9.33	-402.39±5.36	13.393 ±0.023	12.580± 0.022	11.975 ±0.019
J04510093-3402150	04 51 00.93	-34 02 15.04	76.34 ± -8.05	166.66±-8.01	13.541± 0.020	12.826± 0.022	12.294± 0.024
J06244595-4521548	06 24 45.95	-45 21 54.88	-49.71 ± 37.67	392.74±13.81	14.480± 0.026	13.335± 0.027	12.595± 0.024
J06411840-4322329	06 41 18.40	-43 22 32.93	216.60± 15.76	642.33±9.93	13.751± 0.023	12.941± 0.032	12.451± 0.027
J07193188-5051410	07 19 31.88	-50 51 41.06	199.11 ± 20.49	-46.44±13.78	14.094± 0.029	13.282± 0.033	12.773± 0.026
J09221952-8010399	09 22 19.52	-80 10 39.93	3.39 ± 43.54	-66.69±9.36	15.276± 0.053	14.285± 0.033	13.681± 0.046
J10043929-3335189	10 04 39.29	-33 35 18.91	365.99 ± 25.14	-350.34±14.46	14.480± 0.032	13.490± 0.035	12.924± 0.023
J10365305-3441380	10 36 53.06	-34 41 38.09	-32.54 ± 26.06	-445.75±18.42	15.622± 0.046	14.446± 0.034	13.798± 0.042
J11223624-3916054	11 22 36.24	-39 16 05.49	55.74 ± 15.95	-171.34±15.39	15.705± 0.059	14.682± 0.046	13.875± 0.052
J11544223-3400390	11 54 42.23	-34 00 39.06	-158.55 ± 13.23	28.35±13.07	14.195± 0.031	13.331± 0.027	12.851± 0.032
J12073804-3909050	12 07 38.04	-39 09 05.09	-134.78 ± 24.57	51.10±12.76	14.689± 0.038	13.817± 0.026	13.244± 0.037
J13411160-3052505	13 41 11.60	-30 52 50.53	35.45 ± 30.00	-126.20±18.70	14.607± 0.031	13.725± 0.032	13.081± 0.024
J13595510-4034582	13 59 55.10	-40 34 58.27	44.60 ± 17.69	-491.51±14.21	13.645± 0.023	13.034± 0.027	12.566± 0.027
J14252798-3650229	14 25 27.98	-36 50 23.00	-253.11 ± 23.59	-448.55±28.18	13.747± 0.026	12.575± 0.020	11.805± 0.025
J17534518-6559559	17 53 45.18	-65 59 55.91	-50.42 ± 78.51	-329.02±35.82	14.095± 0.025	13.108± 0.026	12.424± 0.027
J19285196-4356256	19 28 51.97	-43 56 25.64	81.24 ± 17.23	-265.82±25.87	15.199± 0.042	14.127± 0.043	13.457± 0.036
J19360187-5502322	19 36 01.88	-55 02 32.22	210.17 ± 32.98	-273.20±21.09	14.486± 0.037	13.628± 0.034	13.046± 0.031
J20414283-3506442	20 41 42.83	-35 06 44.27	56.16 ± 15.79	-118.14±13.02	14.887± 0.031	13.987± 0.020	13.401± 0.036
J21075409-4544064	21 07 54.09	-45 44 06.47	114.58 ± 40.06	-7.85±30.98	14.915± 0.029	13.953± 0.037	13.380± 0.033
J21420580-3101162	21 42 05.80	-31 01 16.29	45.95 ± 6.20	-97.31±5.09	15.844± 0.066	14.767± 0.053	13.965± 0.050
J21501592-7520367	21 50 15.93	-75 20 36.73	869.96 ± 22.08	-277.81±6.55	14.056± 0.026	13.176± 0.031	12.673± 0.029
J21574904-5534420	21 57 49.04	-55 34 42.05	38.93 ± 24.51	-4.58±12.31	14.263± 0.029	13.440± 0.028	13.002± 0.029
J22064498-4217208	22 06 44.98	-42 17 20.89	140.24 ± 10.35	-174.18±4.41	15.555± 0.065	14.447± 0.061	13.609± 0.055
J23312378-4718274	23 31 23.79	-47 18 27.44	72.07 ± 40.02	0.33±28.02	15.659± 0.067	15.510± 0.149	15.389± 0.196

of Gizis, Reid & Monet (1999) and Dobbie et al. (2002) turned up no brown dwarf members. It was hypothesised that being an older open star cluster (625 Myr; Perryman et al. 1998) any low mass members would have evaporated from the cluster through dynamical evolutionary processes. For a cluster of this age $\sim$ 70% of stars and $\sim$ 85% of brown dwarfs are expected to have escaped the cluster (Adams et al. 2002).

Recently however, studies using deep, wide field surveys such as the UKIRT Deep Infrared Sky Survey (Hogan et al. 2008) have unearthed 12 L dwarf candidate members. Bouvier et al. (2008) have also claimed two T dwarf members from a 16 square degree survey of the cluster's centre.

Chereul, Creze & Bienayme (1998) first identified escaped Hyads, and more recently Bannister & Jameson (2007) have identified 7 L and T field dwarfs that belong to the Hyades moving group. Zapatero Osorio et al. (2007) have confirmed that one of these objects (2MASS J1217110-031113) has a radial velocity consistent with being an escaped member of the Hyades cluster, while two (2MASS J0205293-115930 and 2MASS J16241436+0029158) have radial velocities that are only consistent with being moving group members.

In paper I (Jameson et al. 2008a), we reported the discovery of 15 new moving group candidate members. The Hyades moving group has its convergent point situated at α=$6^h29.48^m$, δ=-6°53.4', and the members have a space velocity of 46 km s^{-1} (Madsen, Dravins & Lindgren, 2002).

After using the moving group method, the Cruz et al. (2003) distance and the convergent point from Madsen et al. (2002), we have found 7 new candidate members of the Hyades moving group (Table 5). These objects are plotted on figure 1 as well as the Jameson et al. (2008a); Hogan et al. (2008) and Bannister & Jameson (2007) dwarfs.

Ursa Major

The Ursa Major moving group has been estimated to have an age of between 300 Myr (Soderblom & Mayor 1993) and 500±100 Myr (King et al. 2003). Castellani et al. (2002) found an age of the group to be 400 Myr. The age of 400±100 Myr is adopted in this work. The convergent point of the Ursa Major moving group is located at α=$20^h18^m.83$, δ=$-34°25'.8$ (J2000, Madsen et al. 2002). The selected objects are shown in Figure 2 and their data in Table 6.

In Jameson et al. (2008b) we defined a relationship between M_K

Table 2. 2MASS name, 2MASS RA, 2MASS Dec, $\mu_\alpha \cos\delta$, μ_δ and 2MASS magnitudes for all the L and T dwarfs for which we measured proper motions using WFCAM.

Name	RA	Dec	$\mu_\alpha \cos\delta$	μ_δ	J	H	K_S
2MASS	J2000		mas yr^{-1}				
J00135779-2235200	00 13 57.80	-22 35 20.09	57.74± 23.76	-60.74± 21.38	15.775± 0.064	14.595± 0.062	14.036± 0.050
J00332386-1521309	00 33 23.86	-15 21 30.94	330.31± 16.25	46.33 ±22.26	15.286± 0.055	14.208± 0.051	13.410± 0.038
J00345684-0706013	00 34 56.84	-07 06 01.32	215.35 ±21.32	-138.55± 15.29	15.531± 0.059	14.566± 0.041	13.942± 0.065
J00511078-1544169	00 51 10.79	-15 44 16.91	76.37± 24.08	-4.25± 19.61	15.277± 0.049	14.164± 0.047	13.466± 0.038
J00584253-0651239	00 58 42.53	-06 51 23.94	158.29± 20.18	-105.96 ±15.76	14.311± 0.023	13.444± 0.028	12.904± 0.032
J01311838+3801554	01 31 18.39	38 01 55.48	393.88± 15.45	-21.70± 13.25	14.679± 0.032	13.696± 0.033	13.054± 0.033
J01353586+1205216	01 35 35.86	12 05 21.67	-45.23± 14.82	-441.77± 14.08	14.412± 0.030	13.527± 0.031	12.918± 0.028
J01410321+1804502	01 41 03.22	18 04 50.20	425.05± 17.35	-32.16± 16.50	13.875± 0.022	13.034± 0.024	12.492± 0.026
J01443536-0716142	01 44 35.36	-07 16 14.23	408.28± 17.56	-187.65± 17.99	14.191± 0.023	13.008± 0.027	12.268± 0.021
J01473344+3453112	01 47 33.45	34 53 11.24	47.24± 18.30	-47.19± 14.84	14.946± 0.037	14.162± 0.040	13.574 ±0.037
J02050344+1251422	02 05 03.44	12 51 42.23	388.84± 12.17	-8.71± 12.15	15.679± 0.055	14.449± 0.046	13.671± 0.035
J02073557+1355564	02 07 35.57	13 55 56.45	259.66± 12.71	-159.71± 13.59	15.462± 0.048	14.474± 0.043	13.808± 0.045
J02085499+2500488	02 08 55.00	25 00 48.82	0.71 ±15.49	25.63± 11.047	16.206± 0.092	14.974± 0.080	14.405± 0.069
J02081833+2542533	02 08 18.33	25 42 53.31	386.59± 14.268	-25.14± 16.73	13.989± 0.023	13.107± 0.029	12.588± 0.025
J02082363+2737400	02 08 23.64	27 37 40.09	231.06± 11.62	-95.82± 16.35	15.714± 0.059	14.560± 0.059	13.872± 0.051
J02112827+1410039	02 11 28.28	14 10 03.95	-66.39± 13.83	-27.22± 13.39	16.128± 0.077	15.423 ±0.089	15.009± 0.123
J02132880+4444453	02 13 28.80	44 44 45.36	-42.20± 13.86	-132.39± 13.23	13.494± 0.021	12.757 ±0.023	12.213± 0.021
J02271036-1624479	02 27 10.36	-16 24 47.95	438.00± 18.91	-286.59± 18.81	13.573± 0.020	12.630± 0.022	12.143± 0.029
J02301551+2704061	02 30 15.51	27 04 06.18	198.10 ±15.38	-13.50± 13.63	14.294± 0.024	13.478± 0.028	12.986± 0.021
J02354756-0849198	02 35 47.57	-08 49 19.80	-12.28 ±16.75	21.33± 16.51	15.571± 0.054	14.812± 0.055	14.191± 0.066
J02361794+0048548	02 36 17.94	00 48 54.82	161.33 ±10.10	-176.33± 19.16	16.098± 0.076	15.265± 0.066	14.666± 0.090
J02394245-1735471	02 39 42.46	-17 35 47.20	61.09±13.80	-79.12± 18.79	14.291± 0.029	13.525± 0.034	13.039± 0.030
J02411151-0326587	02 41 11.52	-03 26 58.78	93.43 ±17.00	-19.87± 13.40	15.799± 0.064	14.811± 0.053	14.035± 0.049
J02424355+1607392	02 42 43.55	16 07 39.27	163.62± 14.16	-198.32± 12.77	15.776± 0.052	14.998± 0.053	14.349± 0.057
J02560189+0110467	02 56 01.89	01 10 46.71	66.48± 22.28	-62.38± 22.06	16.212± 0.102	15.696± 0.184	15.216± 0.175
J03090888-1949387	03 09 08.89	-19 49 38.76	218.00± 11.57	-13.33± 11.90	15.752± 0.055	14.656± 0.062	14.057 ±0.065
J03101401-2756452	03 10 14.01	-27 56 45.27	-118.89± 17.61	-46.70± 16.21	15.795± 0.070	14.662± 0.049	13.959± 0.060
J03105986+1648155	03 11 00.02	16 48 15.68	262.83 ± 19.28	9.38± 17.90	16.025± 0.083	14.932± 0.070	14.312 ± 0.067
J03140344+1603056	03 14 03.45	16 03 05.63	-244.73± 13.31	-49.07± 13.91	12.526± 0.021	11.823± 0.035	11.238 ±0.019
J03164512-2848521	03 16 45.13	-28 48 52.17	105.25± 22.15	-80.97±2 12.88	14.578± 0.039	13.772± 0.035	13.114± 0.035
J03202839-0446358	03 20 28.24	-04 46 40.59	-249.91 ± 17.00	-501.77± 12.26	13.259± 0.021	12.535± 0.022	12.134 ± 0.024
J03264225-2102057	03 26 42.26	-21 02 05.77	103.12± 16.14	-129.87± 18.72	16.134 ±0.093	14.793± 0.075	13.922± 0.065
J03281738+0032572	03 28 17.49	00 32 57.57	201.84 ± 22.19	35.50± 17.21	15.988± 0.091	14.975± 0.074	14.161 ± 0.077
J03301774+0000477	03 30 17.74	00 00 47.44	-8.29 ± 11.88	-33.00± 11.08	16.520± 0.111	15.881± 0.130	15.525 ± 0.182
J03504861-0518126	03 50 48.61	-05 18 12.70	19.91± 11.31	-8.04± 15.59	16.327± 0.092	15.525± 0.094	15.125 ±0.134
J03552337+1133437	03 55 23.37	11 33 43.71	223.16± 22.90	-607.16± 22.07	14.050± 0.020	12.530± 0.029	11.526± 0.019
J03554191+2257016	03 55 41.91	22 57 01.68	171.90± 13.14	-21.38± 10.33	16.111± 0.077	15.052± 0.068	14.284± 0.060
J03572110-0641260	03 57 21.10	-06 41 26.04	133.59± 22.58	27.12±5 25.51	15.953± 0.081	15.060± 0.085	14.599± 0.089
J04070885+1514565	04 07 08.85	15 14 56.60	97.97± 16.28	-74.65± 17.40	16.055± 0.091	16.017± 0.208	15.922± 0.261
J04082905-1450334	04 08 29.18	-14 50 34.33	193.43 ± 11.35	-99.07± 17.75	14.222± 0.028	13.337± 0.029	12.817 ± 0.021
J04090950+2104393	04 09 09.51	21 04 39.38	101.13± 15.14	-148.46± 12.25	15.508± 0.053	14.497± 0.045	13.850± 0.046
J04132039-0114248	04 13 20.39	-01 14 24.86	61.22± 15.29	0.08± 18.67	15.303± 0.047	14.657 ±0.033	14.135± 0.059
J04285096-2253227	04 28 51.05	-22 53 21.0	123.44 ± 16.02	183.48± 15.70	13.507 ± 0.020	12.668± 0.026	12.118 ± 0.024
J04390101-2353083	04 39 00.93	-23 53 09.51	-119.29 ± 22.66	-127.25± 15.05	14.408± 0.026	13.409± 0.027	12.816 ± 0.021
J04532647-1751543	04 53 26.51	-17 51 54.35	50.80± 14.38	-4.14± 14.30	15.142± 0.033	14.060± 0.034	13.466 ± 0.034
J05012406-0010452	05 01 24.19	00 10 46.58	192.82± 11.98	-139.86± 12.29	14.982± 0.036	13.713± 0.033	12.963 ± 0.034
J05021345+1442367	05 02 13.45	14 42 36.78	71.55± 13.68	-1.04± 18.53	14.271± 0.022	13.392± 0.019	12.955± 0.028
J05120636-2949540	05 12 06.36	-29 49 53.12	-6.40± 13.41	96.87± 12.28	15.463± 0.055	14.156± 0.047	13.285 ± 0.041
J05184616-2756457	05 18 46.19	-27 56 45.69	37.27± 17.01	10.72± 19.72	15.262± 0.041	14.295± 0.045	13.615 ± 0.039
J05233822-1403022	05 23 38.22	-14 03 02.29	109.87± 17.14	178.88± 21.12	13.084± 0.021	12.220 ±0.020	11.638 ±0.025
J06050196-2342270	06 05 01.96	-23 42 27.01	-66.48± 15.90	117.68± 18.21	14.512± 0.032	13.727 ±0.035	13.145± 0.030
J06262121+0029341	06 26 21.21	00 29 34.14	47.15± 12.94	1.03± 12.78	15.925± 0.093	15.209± 0.120	14.860± 0.123
J06523073+4710348	06 52 30.74	47 10 34.83	-129.78± 14.67	149.52±7 12.78	13.511± 0.020	12.384± 0.023	11.694 ±0.018
J07171626+5705430	07 17 16.27	57 05 43.05	-17.99± 17.87	67.17± 15.13	14.636± 0.029	13.593± 0.028	12.945± 0.025
J07231462+5727081	07 23 14.62	57 27 08.17	61.39± 22.71	-219.85± 12.31	13.970± 0.024	13.156± 0.028	12.613± 0.029
J07400966+3212032	07 40 09.67	32 12 03.24	-31.98± 11.79	-80.79± 13.45	16.191± 0.090	14.862± 0.059	14.222± 0.059
J07414920+2351275	07 41 49.21	23 51 27.51	-250.22± 12.18	-116.21± 13.32	16.158± 0.100	15.838± 0.185	15.847± 99.99
J07420130+2055198	07 42 01.30	20 55 19.88	-318.67± 12.08	-229.57± 13.15	16.193 ±0.090	15.911± 0.181	15.225± 99.99
J07475631+3947329	07 47 56.31	39 47 32.92	55.04± 14.43	-41.72± 13.26	15.076± 0.039	14.163± 0.038	13.724± 0.044
J07525942+4136344	07 52 59.43	41 36 34.49	0.00± 13.39	43.98± 10.46	16.356± 0.130	15.601± 0.160	15.191± 99.99

Table 2. continued

Name 2MASS	RA J2000	Dec J2000	$\mu_\alpha \cos\delta$ mas yr^{-1}	μ_δ mas yr^{-1}	J	H	K_S
J07533217+2917119	07 53 32.17	29 17 11.93	-88.43± 11.08	-85.34± 13.12	15.516± 0.046	14.527± 0.039	13.849± 0.042
J07554795+2212169	07 55 47.95	22 12 16.94	-9.87± 14.88	-226.48± 17.78	15.728± 0.063	15.669± 0.144	15.753± 0.207
J07584037+3247245	07 58 40.37	32 47 24.55	-204.23± 18.01	-316.21± 12.42	14.947± 0.043	14.111± 0.041	13.879± 0.056
J08014056+4628498	08 01 40.56	46 28 49.84	-194.73± 23.54	-331.056± 19.10	16.275± 0.133	15.452± 0.142	14.536± 0.100
J08053189+4812330	08 05 31.89	48 12 33.10	-455.48± 14.62	61.35± 16.32	14.734± 0.034	13.917± 0.040	13.444± 0.040
J08155674+4524119	08 15 56.75	45 24 11.93	-31.49± 15.56	-42.96±13.18	16.057± 0.078	15.233± 0.093	14.812± 0.097
J08202996+4500315	08 20 29.96	45 00 31.52	-104.16± 25.82	-299.16±14.76	16.279± 0.107	15.000± 0.086	14.218± 0.065
J08234818+2428577	08 23 48.18	24 28 57.71	-160.30± 19.81	73.19± 19.17	14.986± 0.042	14.060± 0.044	13.377± 0.029
J08290664+1456225	08 29 06.64	14 56 22.56	-49.41± 16.67	-227.11± 18.47	14.750± 0.028	13.801± 0.035	13.166± 0.031
J08304878+0128311	08 30 48.78	01 28 31.15	222.11± 15.89	-310.54± 15.36	16.289± 0.111	16.140± 0.213	16.358± 99.99
J08320451-0128360	08 32 04.52	-01 28 36.05	63.83± 13.19	26.73± 15.10	14.128± 0.028	13.318± 0.022	12.712± 0.026
J08355829+0548308	08 35 58.30	05 48 30.85	-109.70± 12.79	-15.00± 15.54	14.533± 0.034	13.683± 0.036	13.168± 0.033
J08472872-1532372	08 47 28.73	-15 32 37.21	149.07± 16.12	-177.78 ±15.82	13.513± 0.023	12.629± 0.026	12.061± 0.021
J08523490+4720359	08 52 34.91	47 20 35.91	-28.69 ±18.13	-386.53± 13.88	16.182± 0.108	15.419± 0.146	14.718± 0.116
J08564793+2235182	08 56 47.94	22 35 18.21	-184.24± 19.74	3.01± 20.71	15.679 ±0.064	14.580± 0.052	13.951± 0.046
J08575849+5708514	08 57 58.49	57 08 51.42	-413.61± 20.52	-353.43±16.85	15.038± 0.038	13.790± 0.041	12.962 ±0.028
J08592547-1949268	08 59 25.48	-19 49 26.89	-310.77 ±13.51	-78.59±15.95	15.527± 0.052	14.436± 0.041	13.751± 0.057
J09095749-0658186	09 09 57.49	-06 58 18.64	-174.53± 14.89	39.70± 16.48	13.890± 0.021	13.090± 0.020	12.539± 0.024
J09153413+0422045	09 15 34.14	04 22 04.59	-88.46± 18.02	40.27± 20.60	14.548 ±0.028	13.531± 0.031	13.011± 0.041
J09183815+2134058	09 18 38.16	21 34 05.82	353.41± 14.50	-454.02± 15.66	15.662± 0.060	14.580± 99.99	13.903± 0.042
J09201223+3517429	09 20 12.23	35 17 42.97	-172.34± 13.35	-185.26± 12.72	15.625± 0.062	14.673± 0.056	13.979± 0.061
J09211410-2104446	09 21 14.11	-21 04 44.60	260.71± 15.59	-900.91± 13.20	12.779± 0.021	12.152± 0.020	11.690± 0.021
J09283972-1603128	09 28 39.72	-16 03 12.86	-132.85±15.59	36.00± 13.20	15.322 ±0.041	14.292± 0.036	13.615± 0.050
J09352803-2934596	09 35 28.04	-29 34 59.62	8.84± 18.10	86.92±14.61	14.036± 0.026	13.312± 0.027	12.822± 0.026
J10185879-2909535	10 18 58.79	-29 09 53.56	-323.01± 21.04	-82.84± 15.20	14.213± 0.028	13.418± 0.022	12.796± 0.021
J10432508+0001482	10 43 25.00	00 01 46.92	-159.31± 15.56	-143.85± 21.27	15.935 ± 0.080	15.208 ± 0.072	14.472 ± 0.100
J12285538+0050440	12 28 55.37	00 50 43.94	-32.33±3.54	-12.41±2.06	15.613± 0.060	14.825± 0.059	14.162 ± 0.077
J16360078-0034525	16 36 00.59	00 34 54.36	-344.71± 14.13	-200.89± 14.86	14.590 ± 0.043	13.904± 0.042	13.415 ± 0.035
J17434148+2127069	17 43 41.59	21 27 09.16	165.91± 14.890	248.68± 14.47	15.830± 0.088	14.785± 0.064	14.321 ± 0.097

Table 3. Name, μ from Schmidt et al. (2007), θ from Schmidt et al. (2007), μ, θ for all the L and T dwarfs common with Schmidt et al. (2007).

Name 2MASS	μ (Schmidt) " yr^{-1}	θ (Schmidt) $^\circ$	μ " yr^{-1}	θ $^\circ$
J0213+44	0.17±0.06	195±21	0.138±0.01	197±6
J0255-47	1.23±0.11	120±5	1.18±0.01	117±1
J0314+16	0.25±0.07	254±13	0.25±0.01	256±3
J0355+11	0.70±0.05	159±3	0.64±0.02	160±2
J0439-23	0.20±0.17	220±37	0.17±0.02	223±6
J0445-30	0.38±0.11	161±23	0.43±0.01	160±1
J0523-14	0.08±0.16	21±105	0.2±0.02	32±5
J0624-45	0.28±0.19	19±52	0.39±0.01	352±5
J0651+47	0.14±0.05	336±13	0.19±0.013	319±4
J0847-15	0.27±0.05	146±10	0.23±0.015	140±4
J0915+04	0.14±0.15	284±84	0.097±0.018	294±11
J0921-21	0.98±0.15	163±10	0.93±0.113	164±1
J1425-36	0.53±0.10	204±7	0.52±0.03	209±3
J1753-65	0.36±0.09	178±24	0.33±0.04	189±13

and $J-K$ for the L dwarf sequence in the Hyades and Ursa Major moving groups and the Pleiades, Upper Scorpius and Alpha Persei open star clusters. This allowed a photometric relationship for estimating the age of any field dwarf to be derived.

Using the new objects presented here and in Paper I (Jameson et al. 2008a) we are able to refine the relationship derived in (Jameson et al. 2008b) for the Ursa Major moving group. Previously, this relationship (which takes the form $J-K$=m $\times M_K$+c, in the MKO system) had a gradient (m)=2.88 and c=11.22. This was only based on the 2 brown dwarfs in the L dwarf sequence from Bannister & Jameson (2007), and was obviously incorrect. The sequence shown in Figure 2 of Jameson et al. (2008b) crossed that for the Hyades, and the gradient of 2.88, did not fit, with the gradients of the other clusters, which had approximately parallel sequences, all with gradients of $\sim$1.98, the value that was suggested by the authors as being more appropriate. Using our new L dwarfs,

Table 4. Name, μ from Caballero (2007), θ from Caballero (2007), μ, θ for all the L and T dwarfs common with Caballero (2007).

Name 2MASS	$\mu_\alpha \cos\delta$ (Caballero)	μ_δ(Caballero)	$\mu_\alpha \cos\delta$	μ_δ
		mas yr^{-1}		
J0255-47	1060±50	-630±50	1052.88 ± 11.17	-546.54±6.17
J0445-30	167±12	-424±12	158.07 ± 9.33	-402.39±5.36
J0451-34	94±17	114±16	76.34 ± -8.05	166.66±-8.01
J0719-50	140±30	-10±30	199.11 ± 20.49	-46.44±13.78
J0847-15	-130±160	-20±180	149.07± 16.12	-177.78 ±15.82
J0909-07	-280±190	110±180	-174.53± 14.89	39.70± 16.48
J0921-21	100±60	-900±60	260.71± 15.59	-900.91± 13.20



Figure 1. M_K, J-K colour magnitude diagram for the Hyades moving group. The cluster members members identified by Hogan et al. (2008) are plotted as filled squares. All of the objects that were previously selected as moving group members are marked as filled diamonds (Jameson et al. 2008a; Bannister & Jameson 2007). The selected members are marked by a box for objects from the SofI data and by circles for the WFCAM objects. The errors are poissonion and from the photometry only. The solid and dashed lines represent the single star (solid line) and binary (dotted line) cluster sequences as shown in Hogan et al. (2008).

Table 5. Name, spectral type, J, H, K magnitudes, $\Delta\theta$, d_{mg}/d_{sp} and d_{sp} for the potential Hyades moving group members discussed.Where two spectral types have been given, they are in the order of optical spectral type, infrared spectral type.

Name	spt	J	H	K	$\Delta\theta$	d_{sp}	d_{mg}/d_{sp}
					$^\circ$	pc	pc
J0131+38	L4	14.68± 0.03	13.70± 0.03	13.05± 0.03	12.18±1.93	93.15	1.19
J0141+18	L1/L4	13.88± 0.03	13.03 ±0.02	12.49± 0.03	12.65±2.21	94.33	0.96
J0205+12	L5	15.68± 0.06	14.45± 0.05	13.67± 0.04	9.19±1.79	91.28	0.93
J02081+25	L1	13.99± 0.03	13.11± 0.03	12.59± 0.03	12.19±2.47	93.72	1.01
J0357-06	L0	15.95± 0.08	15.06± 0.09	14.60± 0.09	5.01±10.67	78.52	1.02
J0624-45	L5	14.48± 0.03	13.34± 0.03	12.60± 0.02	8.69± 5.41	15.41	1.27
J1928-43	L5	15.20± 0.04	14.13± 0.04	13.46± 0.04	6.70± 3.74	21.46	1.04

and those found by Jameson et al. (2008a), we used the colour conversions of Stephens & Leggett (2004) to recalculate the relationship on the MKO system. Our new values are m=2.0746±0.281, c=7.92052±0.4481, which is more in keeping with the values for the other clusters. The relationship in the 2MASS colour system is plotted on Figure 2, as well as the sequence as defined by the Bannister & Jameson (2007) dwarfs.

Pleiades

The Pleiades cluster is 125 Myr old and is situated at a distance of 130 pc (Stauffer, Schultz & Kirkpatrick 1998). As a cluster it has been studied in depth and has been found to contain many brown dwarfs (Casewell et al. 2007; Lodieu et al. 2007b; Bihain et al. 2006; Moraux et al. 2003). The Pleiades moving group has a convergent point of $85.04^\circ \pm 3.67$, $-39.11^\circ \pm 6.92$ (Madsen et al. 2002). This convergent point is very close to that of many other moving groups such as Alpha Persei ($96^\circ.78 \pm 1.96$, $-23^\circ.27 \pm 3.67$; Madsen et al. 2002, 50 Myr; Lyngå, G 1987), Tucana/Horologium (30 Myr; Zuckerman & Song 2004) and the AB Dor moving group (50 Myr; Zuckerman & Song 2004) (see Zuckerman & Song 2004 for a review), and it has been theorised that many of these moving groups have a common origin (Ortega et al. 2007). Thirteen new candidate members were found using the moving group method. 6 objects with declinations $<-30^\circ$ and 7 objects with declinations $>-30^\circ$ (Figure 3, Table 7). As with the Hyades and Ursa Major clusters, the Pleiades is old enough to expect that some mass segregation has occurred, and thus it is not unreasonable to search the whole sky for members of the moving group. This is not the case however for many of the younger, southern moving groups.

The new moving group members are detailed in Table 7.

Other moving groups

Many of the southern moving groups have similar convergent points and velocities. To determine if any of our southern dwarfs are members of these moving groups, we have used the moving group method as for the Hyades, Ursa Major and Pleiades moving groups, but have then used the isochrone for Upper Scorpius as developed by Jameson et al. (2008b). This can be then used as an age indicator for the younger clusters. If the selected dwarfs fall on or near the isochrone, then they are young (age $<$10 Myr) and are considered candidate members. Radial velocity measurements are needed to confirm the membership of these objects however.

Simply by using the moving group method, we have found $<$10 candidate members of the TW Hydra, Tucana/Horlogium, Beta Pictoris, AB Doradus and η Chamaeleon moving groups. Because these clusters all have similar convergent points and velocities (Zuckerman & Song 2004) some of the candidate members are found in more than one cluster.

When plotted on the M_K, $J-K$ colour magnitude diagram, using Stephens & Leggett (2004) to convert the colours into the MKO system, with the Upper Scorpius and Alpha Perseus cluster isochrones from Jameson et al. (2008b), all of the candidate members sit lower then Alpha Per, indicating that these dwarfs cannot be moving group members at the calculated distances.

CONCLUSIONS

This paper continues the work presented in Jameson et al. (2008a), which presented proper motions of 143 L and T dwarfs. This paper presents a further 126 proper motions. Thus the large majority of field L and T dwarfs discovered by 2MASS, DENIS and the SDSS now have known proper motions. From these data we find a further 2 wide binary L dwarfs, both with M dwarf companions. A further 3 high velocity L dwarfs have been discovered, which we assume are old thick disc L dwarfs. Finally, we find 7 more potential members of the Hyades moving group, 4 members of the Ursa Major moving group and 13 potential members of the Pleiades moving group. We have found no members of the young southern moving groups, TW Hydra, Tucana/Horologium, Beta Pictoris, AB Doradus or η Chamaeleon. We have also used these new members of the Ursa Major moving group to refine the L dwarf sequence for the group that was defined by Jameson et al. (2008b).

ACKNOWLEDGEMENTS

SLC was supported by STFC for the duration of this work. MBU is supported by a STFC Advanced Fellowship. Observations were made at the United Kingdom Infrared Telescope, which is operated by the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council.Observations were also made a the New Technology Telescope which is operated by ESO. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. Research has benefited from the M, L, and T dwarf compendium housed at DwarfArchives.org and maintained by Chris Gelino, Davy Kirkpatrick, and Adam Burgasser. This research has made use of NASA's Astrophysics Data System Bibliographic Services.

REFERENCES

Adams T., Davies M. B., Jameson R. F., Scally A., 2002, MNRAS, 333, 547

Allan A, Jenness T, Economou, F, Currie, M. J., Bly M. J., 2002, in ASP Conf. Ser., Vol. 281 Astronomical Data Analysis Software and Systems XI, eds Edited by David A. Bohlender, Daniel Durand, and Thomas H. Handley (San Francisco: ASP), 311

Bannister N. P., Jameson R. F., 2007, MNRAS, 378, L24

Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1998, A&A, 332, 403

Bihain G., Rebolo R., Béjar V. J. S., Caballero J. A., Bailer-Jones C. A. L., Mundt R., Acosta-Pulido J. A., Manchado Torres A., 2006, A&A, 458, 805

Bouvier J., et al., 2008, A&A, 481, 661

Burgasser A. J., Liu M. C., Ireland M. J., Cruz K. L., Dupuy T. J., 2008, ApJ, 2008arXiv0803.0295B

Burrows A., Hubbard W. B., Lunine J. I., Liebert J., 2001, RvMP, 73, 719

Caballero J. A., 2007, ApJ, 667, 520

Casali M., et al., 2007, A&A, 467, 777

Casewell S. L., Dobbie P. D., Hodgkin S. T., Moraux E., Jameson R. F., Hambly N. C., Irwin J., Lodieu N., 2007, MNRAS, 378, 1131

Castellani V., Degl'Innocenti S., Prada Moroni P. G., Tordiglione V., 2002, MNRAS, 334, 193



Figure 2. M_K, J-K colour magnitude diagram for the Ursa Major moving group. The cluster members members identified by Jameson et al. (2008a) and Bannister & Jameson (2007) are marked by filled diamonds. The selected members are marked by a box for objects from the SofI data and by filled circles for the WFCAM objects. The errors are poissonion and from the photometry only. The solid line indicates a possible single star sequence from Bannister & Jameson (2007). The straight line fit to the new L dwarf sequence is shown by a dotted line and the dashed line represents what we believe to be the turnaround of the sequence.

Table 6. Name, spectral type, J, H, K magnitudes, $\Delta\theta$, d_{mg}/d_{sp} and d_{sp} for the potential Ursa Major moving group members discussed.Where two spectral types have been given, they are in the order of optical spectral type, infrared spectral type.J0920+35 is a binary (Reid et al. 2001), discovered using HST. Is suspected to be an equal mass binary.

Name	spt	J	H	K	$\Delta\theta$ $^\circ$	d_{sp} pc	$_{mg}/d_{sp}$
J0218-31	L3	14.73±0.04	13.81±0.04	13.15±0.03	6.29± 5.14	25.85	0.86
J0310-27	L5	15.80± 0.07	14.66± 0.05	13.96± 0.06	1.48 ±7.36	28.24	0.80
J0815+45	L1	16.06± 0.08	15.23± 0.09	14.81± 0.10	12.04± 15.89	64.88	1.09
J0920+35	L6.5/T0	15.63± 0.06	14.67± 0.06	13.98± 0.06	5.30± 2.96	19.06	0.79

Table 7. Name, spectral type, J, H, K magnitudes, $\Delta\theta$, d_{mg}/d_{sp} and d_{sp} for the potential Pleiades moving group members discussed.Where two spectral types have been given, they are in the order of optical spectral type, infrared spectral type.

Name	spt	J	H	K	$\Delta\theta$ $^\circ$	d_{sp} pc	d_{mg}/d_{sp}
J0032-44	L0	14.78± 0.03	13.86± 0.03	13.27± 0.04	13.71± 2.51	40.70	0.86
J0058-06	L0	14.31± 0.03	13.44± 0.03	12.90 ±0.03	4.83± 5.19	32.85	0.99
J0117-34	L2	15.18± 0.03	14.20± 0.04	13.49± 0.04	4.84± 4.28	37.58	1.18
J0125-34	L2	15.52± 0.05	14.47± 0.05	13.90± 0.05	18.82± 10.14	44.03	0.90
J0144-07	L5	14.20± 0.03	13.01± 0.03	12.27± 0.02	13.87± 2.28	13.49	1.02
J0208+27	L5	15.71± 0.06	14.56± 0.06	13.87± 0.05	15.56± 3.60	27.21	0.903
J0236+00	L6/L6.5	16.10± 0.08	15.27± 0.07	14.67 ±0.09	8.13± 3.58	26.22	1.00
J0316-28	L0	14.57± 0.04	13.77± 0.03	13.11± 0.03	4.55± 7.30	37.02	1.17
J0357-44	L0	14.37 ±0.03	13.53± 0.03	12.91± 0.03	4.12± 6.85	33.71	0.99
J0409+21	L3	15.51± 0.05	14.50± 0.05	13.85± 0.05	16.18± 4.55	37.03	0.94
J1425-36	L5	13.74 ±0.03	12.57± 0.02	11.81± 0.03	6.23± 2.76	11.00	1.11
J1928-43	L5	15.20 ±0.04	14.13± 0.04	13.46± 0.04	3.80± 3.74	21.46	1.05
J1936-55	L5	14.49 ±0.04	13.63± 0.03	13.05± 0.03	15.30± 4.84	15.46	1.17



Figure 3. M_K, *J-K* colour magnitude diagram for the Pleiades moving group. The cluster members members identified by Casewell et al. (2007), Lodieu et al. (2007b), Moraux et al. (2003) and Bihain et al. (2006) are plotted as filled squares. All of the objects that were selected as moving group members are marked as filled diamonds (Jameson et al. 2008a). The selected members are marked by a box for objects from the SofI data and by filled circles for the WFCAM objects. The errors are poissonion and from the photometry only.

Chabrier G., Baraffe I., Allard F., Hauschildt P. H., 2000, ApJ, 542, 464

Chereul E., Creze M., Bienayme O., 1998, A&A, 340, 384

Costa E., Mendez R. A., Jao W.-C., Henry T. J., Subasavage J. P., Ianna P. A., 2006, AJ, 132, 1234

Cruz K. L., Reid I. N., Liebert J., Kirkpatrick J. D., Lowrance P. J., 2003, AJ, 126, 2421

Currie, M. J. 2003, in ASP Conf. Ser., Vol. 314 Astronomical Data Analysis Software and Systems XIII, eds. F. Ochsenbein, M. Allen, and D. Egret (San Francisco: ASP), 460

Dehnen W., 1998, AJ, 115, 2384

Delfosse X.,Tinney C. G., Forveille T., Epchtein N., Borsenberger J., Fouqué P., Kimeswenger S., Tiphène D., 1999, A&AS, 135,41

DENIS Consortium, 2005, The DENIS database, VizieR Online Data Catalog

Dobbie P. D., Kenyon, F., Jameson, R. F., Hodgkin, S. T., Hambly, N. C., Hawkins, M. R. S., 2002, MNRAS, 329, 543

Dye S., et al., 2006, MNRAS, 372, 1227

Gizis J. E., Reid I. N., Monet D. G., 1999, AJ, 118, 997

Hambly N. C. et al., 2001a, MNRAS, 326, 1279

Hambly N. C., Davenhall A. C., Irwin M. J., MacGillivray H. T., 2001b, MNRAS, 326, 1315

Hawley S. L., et al., 2002, AJ, 123, 3049

Hogan E., Jameson R. F., Casewell S. L., Osbourne S. L., Hambly N. C., 2008, MNRAS, 2008arXiv0805.1189H

Kendall T. R., et al., 2007, A&A, 466, 1059

King J. R., Villarreal A. R., Soderblom D. R., Culliver A. F., Adelman S. J., 2003, AJ, 125, 1980

Jameson R. F., Casewell S. L., Bannister N. P., Lodieu N., Keresztes K., Dobbie P. D., Hodgkin S. T. 2008a, MNRAS, 384, 1399

Jameson R. F., Lodieu N., Casewell S. L., Bannister N. P., Dobbie P. D., 2008b, MNRAS, 385, 1771

Lawrence A., et al., 2007, MNRAS, 379, 1599

Leggett S. K., et al., 2002, ApJ, 564, 452

Lodieu N. et al., 2007a, MNRAS, 379, 1423

Lodieu N., Hambly N. C., Jameson R. F., Hodgkin S. T., Carraro G., Kendall T. R., 2007b, MNRAS, 374, 372

Lyngå, G, 1987, Catalogue of open cluster data, VizierR Online Data Catalog

Madsen S., Dravins D, Lindgren L., A&A, 2002, 381, 446

Moorwood, A., Cuby J. G., Lidman C., 1998, The Messenger 91, 9

Moraux E., Bouvier J., Stauffer J. R.,Cullindre J-C., 2003, A&A, 400, 891

Ortega V. G., Jilinski E., de La Reza R., Bazzanella B., 2007, MNRAS, 377, 441

Perryman M. A. C., et al., 1998, A&A, 331, 81

Reid I. N., Lewitus E., Burgasser A. J., Cruz K. L., 2006, ApJ, 639, 1114

Reid I. N., Gizis J. E., Kirkpatrick J. D., Koerner D. W., 2001, AJ, 121, 489

Schmidt S. J., Cruz K. L., Bongiorno B. J., Liebert J., Reid I. N., 2007, AJ, 133, 2258

Skrutskie M. F., et al., 2006, AJ, 131, 1163

Soderblom D. R., Mayor M., 1993, AJ, 105, 226

Stauffer J. R., Schultz G., Kirkpatrick J. D., 1998, ApJ, 499, 199L

Stephens D. C., Leggett S. K.,2004, PASP, 116, 9

Warren S. J., et al., 2007, MNRAS, 375, 213

Wei J. Y., Xu D. W., Dong X. Y., Hu J. Y., 1999, A&AS, 139, 575

York D. G., et al., 2000, AJ, 120, 1579

Zapatero Osorio M. R., Martín E. L., Béjar V. J. S., Bouy H., Seshpande R., Wainscoat R. J., 2007, ApJ, 666, 1205

Zuckerman B., Song I., 2004, ARA&A, 42, 685